

The UK Vaccination Schedule

- At birth:** BCG (Tuberculosis) for babies with infected or previously-infected family members; or from UK areas where incidence of TB is 40 in 100,000 or greater; or whose parents or grandparents were born in a high-risk country; or immigrants from high-risk countries.
- Hepatitis B for babies born to mothers who are chronic carriers of the Hepatitis B virus or who have had acute Hepatitis B during pregnancy.
- At 2 mths:** Diphtheria, Whooping Cough, Tetanus, Polio, Hib [Pediace® 5-in-1] plus Pneumococcal Vaccine [Prevna®] (i.e. 2 injections)
- At 3 mths:** Pediace® plus Meningitis C [Meningitec®] (i.e. 2 injections)
- At 4 mths:** Pediace®, Meningitec® and Prevna® (i.e. 3 injections)
- NB: 19 vaccines by age 5 months*
- At 12 mths:** Meningitis C plus Hib [Menitorix®] (in combination)
- At 15 mths:** Measles, Mumps, Rubella (=MMR); plus Prevna® (i.e. 2 injections)
- NB: 25 vaccines by age 16 months*
- At 3 yrs 4 mths to 5 yrs:** Diphtheria, Whooping Cough, Tetanus, Polio [Infanrix/IPV®] DTaP IPV; plus MMR (i.e. 2 injections) plus Hepatitis B booster for babies immunised at birth (i.e. born to mothers infected with Hepatitis B)
- NB: 32 vaccines by age 5 years*
- At 12-13 yrs:** Human Papilloma Virus [Cervarix®*] for girls aged 12 and upwards (3 injections over 6 months); with catch-up programmes planned for 2009-10 for girls aged 15-18 years
- At 15 yrs:** Diphtheria, Tetanus, Polio [Revaxis® dT IPV]; plus MMR if fewer than 2 previous doses (i.e. 2 injections)

References:

- <http://www.vaccineriskawareness.com>
 - <http://www.immunisation.nhs.uk/immunisation-schedule>
-